



Proceeding Paper

Application of Dairy Manure Amended with Mineral Fertilizer on Stubble-Covered Soil: Effects on Ammonia Emissions [†]

Arejacy A. Silva ^{1,2} and David Fangueiro ^{2,*}

¹ Instituto Federal de Educação, Ciência e Tecnologia de São Paulo, Av. Prof. Celso F. Silva, 1333, Avare 18707-150, Brazil; arejacy.silva@ifsp.edu.br

² LEAF, Instituto Superior de Agronomia, Universidade de Lisboa, Tapada da Ajuda, 1349-017 Lisboa, Portugal

* Correspondence: dfangueiro@isa.ulisboa.pt

[†] Presented at the 1st International Electronic Conference on Agronomy, 3–17 May 2021; Available online: <https://sciforum.net/conference/IECAG2021>.

Abstract: The reduction in the manure application rates through enrichment with mineral fertilizer has the potential to reduce costs, decrease environmental pollution, and extend the manure benefits to greater acreage. A pot experiment was carried out to assess ammonia emissions from dairy manure amended with mineral fertilizers applied on wheat stubble. The treatments were: control (no fertilization), urea (U), calcium ammonium nitrate (AN), dairy manure (MAN), urea + dairy manure (UMAN), and calcium ammonium nitrate + dairy manure (ANMAN). A dynamic chamber system was used to measure NH₃ emissions during seven days after soil application. UMAN and ANMAN treatments led to higher NH₃ emissions than each isolated component. This might be motivated by the manure pH. Thus, the enrichment of dairy manure with U or AN for application on stubble-covered soil should not be recommended. Nevertheless, some manure pre-treatments, such as acidification, or the use of other mineral fertilizers might improve such solution.

Keywords: ammonia; nitrogen; organic-mineral fertilizer; manure; no-tillage



Citation: Silva, A.A.; Fangueiro, D. Application of Dairy Manure Amended with Mineral Fertilizer on Stubble-Covered Soil: Effects on Ammonia Emissions. *Biol. Life Sci. Forum* **2021**, *3*, 19. <https://doi.org/10.3390/IECAG2021-10017>

Published: 11 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Mineral fertilizers are the major source of nitrogen (N) to crops [1,2]. However, manure application contributes to the increase in soil organic matter and the addition of beneficial microbes, besides also delivering nutrients to plants [3]. The transport costs within and between farms and higher application rates needed relative to mineral fertilizers are some of the limitations associated with manure application [4].

The mixture of manure and mineral fertilizer might be an alternative to decrease manure application rates, enabling the coverage of more agricultural lands, improving application efficiency, and reducing over-fertilization with phosphorus, as generally occurs when using manure [5]. The chances of water contamination through manure runoff following heavy precipitation or snow melting can be reduced by applying manure on soil covered with crop residues [6]. However, livestock manure is the main contributor to ammonia (NH₃) emissions. The ammonia emitted from livestock manure and mineral fertilizers constitutes an important loss of reactive nitrogen and represents a threat to human health. Furthermore, the NH₃-N carried to land or water may surpass the critical nitrogen load of the ecosystems, causing eutrophication [7–9].

Despite the evident benefits of the joint application of manure and mineral fertilizer, it is necessary to evaluate all potential impacts of this technique on the environment and crop production. The present work aimed to assess the impact of urea and calcium ammonium nitrate amendment to manure, right before application to stubble-covered soil, on ammonia emissions.

2. Materials and Methods

The experiment was carried out for seven days in a greenhouse at Instituto Superior de Agronomia, Lisbon, Portugal (coordinates 38°42'29.786" N; 9°11'6.18" W). The average maximum and minimum temperatures in the greenhouse during the experimental period were 26.4 °C and 10.6 °C. The manure used came from the storage tank of a typical commercial dairy farm located in the Setubal region, Portugal.

The pots were filled with 2 kg of clay loam soil referenced as Vertic soil (texture characteristics: 16.8% coarse sand, 33.7% fine sand, 20.9% silt, and 28.6% clay), covered by wheat stubbles (300 g m⁻²). The treatments, with 3 replicates, conducted in a completely randomized design, were unfertilized soil (control) and soil application of: urea (U), calcium ammonium nitrate (AN), dairy manure (MAN), urea + dairy manure (UMAN), and calcium ammonium nitrate + dairy manure (ANMAN). The manure, mineral fertilizers, and their combinations were spread by hand on the stubbles at a rate of 0.5 g of total-nitrogen (Total-N) per pot. For the organic–mineral combinations, the mixture was prepared right before the application and each component contributed 50% of total-N. Soil moisture was initially set up at 60% of soil water holding capacity, before fertilizer application, and then adjusted only on the fourth day by water addition. The urea (U) used in this experiment contained 46% of total-N and the calcium ammonium nitrate (AN) contained 27% of total-N plus 4% of CaO. The main soil and dairy cattle manure characteristics are shown in Table 1.

Table 1. Main soil and manure characteristics (mean of 3 replicates).

Parameters	Soil	Manure
Total Nitrogen (g kg ⁻¹)	1.70	11.5 **
NH ₄ -Nitrogen (g kg ⁻¹)	0.01	3.86 **
pH (H ₂ O)	7.1	7.40
Dry Matter * (%)	-	28.50

* % of raw manure. ** Based on dry matter.

2.1. Ammonia Measurements

After the application of the fertilizers on the stubble, the ammonia emissions were collected from each pot topped with a PVC chamber (0.035 m² area), through a dynamic chamber system, similar to the one described by [10], for 7 days. A constant airflow (3 L min⁻¹) was maintained inside the chambers, using a laboratory suction pump regulated by a needle valve, and an acid trap containing 200 mL of H₃PO₄ (0.05 M), connected to each chamber to collect the NH₃ emitted. The acid solution in each acid trap was replaced after 4, 8, and 12 h, in the first 24 h, twice a day in the second and third days, and then every 24 h until day 7. At the end of each sampling period, the total ammoniacal N (TAN) content in the solution of each acid trap was analyzed by automated segmented-flow spectrophotometry [11]. Ammonia emission rates (E , mg N m⁻² h⁻¹) for each sampling period were calculated according to Equation (1).

$$E = \frac{\text{TAN} \times V}{S \times t} \quad (1)$$

where TAN is the total ammoniacal N concentration of the acid solution (in mg L⁻¹), V is the volume of acid solution (in L), S is the soil surface area (in m²), and t is the time interval of the NH₃ trapping (in h). Total NH₃ emissions were also expressed as the sum of the amount of NH₃ emitted during each time interval and expressed as % of total nitrogen applied.

2.2. Statistical Analysis

Statistical analysis of ammonia emissions, using ANOVA, was undertaken with Statistix 9. The least significant differences (LSD) were used to compare means with a probability level of 5%, and normalization of the data was not required.

3. Results

The cumulative ammonia emissions expressed in mg NH₃-N per pot and as % of Total-N applied are presented in Table 2. The highest cumulative amounts of volatilized ammonia ($p < 0.05$) were registered when manure was amended with both mineral fertilizers.

Table 2. Cumulative ammonia emission expressed as mg NH₃-N pot⁻¹ and as a percentage of total nitrogen applied (removed the control emission). For each parameter, in the same column, values followed by different letters are significantly different based on the LSD test (mean of three replicates).

	Cumulative NH ₃ Emission mg NH ₃ -N pot ⁻¹	% of Total-N Applied
UMAN	117.5 ^a	23.4 ^a
ANMAN	96.8 ^a	19.3 ^a
MAN	62.7 ^b	12.5 ^b
U	25.7 ^c	5.0 ^c
AN	5.9 ^{cd}	1.1 ^c
Control	0.51 ^d	-

In MAN, 12.5% of total-N applied (~37% of TAN applied) was lost as ammonia. Despite the relatively low total-N loss, this represents an amount significantly higher than those observed in mineral fertilizers (U and AN), but less than N lost from UMAN (23.4%) and ANMAN (19.3%).

The daily ammonia emission rates showed that the highest peak was reported in UMAN followed by ANMAN. Daily rates from all fertilizers peaked on the first day and then decreased to negligible levels after the fourth day, except U, which only reached the peak on the third day (Figure 1a,b), and at the end of the seventh day. Albeit in a small amount, urea still emitted more ammonia than the other treatments. All the treatments recorded another slight peak after the complementary irrigation performed on the fourth day.




Figure 1. NH₃-N daily emission rates (mg pot⁻¹ h⁻¹). Error bars represent the standard error values (mean of three replicates). (a) EmisScheme 2. NH₃-N emission dynamics (percentage of the total NH₃-N emitted). (a) NH₃-N emissions dynamics from UMAN, MAN, and ANMAN. (b) NH₃-N emissions dynamics from AN, U, and control.

The dynamics of emissions, presented as the percentage of NH₃-N daily emission rates related to the total NH₃-N emissions (100%), are shown in Figure 2a,b. More than 80% of total NH₃-N emissions from UMAN and ANMAN occurred in the two first days, while MAN reached that mark only on the third day. Diversely, the ammonia emissions from AN, U, and control were much less intense, highlighting urea that only reaches half of its total NH₃-N emission on the fourth day.




Figure 2. NH₃-N emission dynamics (percentage of the total NH₃-N emitted). (a) NH₃-N emissions dynamics from UMAN, MAN, ANMAN. (b) NH₃-N emissions dynamics from AN, U, Control.

4. Discussion

It is noteworthy that the joint application of manure and urea or manure and calcium ammonium nitrate led to higher NH₃-N emission than the sum of emissions from each of the component separately (Table 2). UMAN emitted 1.87 and 4.47 times more NH₃-N than MAN and U, respectively. ANMAN emitted 1.54 and 16.4 times more than MAN and AN, respectively.

The NH₃-N losses from manure were expected to be higher than from mineral fertilizers [12] although, according to [13], losses of ammonia from solid manure applied to the soil are not well understood as the emissions from slurry application. Thus, the cumulative ammonia emission, as well as the N lost relative to the applied total-N, from MAN is probably justified by its low NH₄-N content and the susceptibility to the formation of surface crust due to its high dry matter level [7,14].

The highest NH₃-N emission from manure amended with mineral fertilizers might be explained by the manure pH, which favored the dominance of ammonia over ammonium as described by [15,16]. In addition, we can hypothesize that the highest and more intense ammonia emissions (Figure 1a,b and Figure 2a,b) from UMAN might be determined firstly by the contact of the urea with the urease from manure in a medium wetter than the soil, favoring the conversion from urea to ammoniacal nitrogen, and thereafter the NH₃ volatilization stimulated by the alkaline pH. The low ammonia emission from calcium ammonium nitrate, which contains 50% of the nitrogen in nitric form, is in agreement with data reported in other studies [8,17]. The N loss from U was below expectations, even if in agreement with the values described by [18]. Thereby, NH₃-N emission from U was probably kept at a low level as, after dissolution, the fertilizer was protected from airflow and solar radiation by the stubbles, as already observed by [19] after application of pig slurry on stubble, these authors attributed the reduction in ammonia emission to the protection provided by the stubble layer. Additionally, an initial lag phase was observed in the NH₃-N daily emission rates from U (Figure 1b), influencing its dynamics (Figure 2b), probably due to the time needed to convert urea into ammonia, through the urease enzyme [20,21]. Thus, assessment of ammonia emissions from urea applied to stubble-covered soil should last for more than the seven days usually considered in studies dealing with ammonia emissions from manure.

5. Conclusions

The application of dairy manure mixed with urea or calcium ammonium nitrate on stubble-covered soil stimulates ammonia emissions relative to the isolated application of manure or mineral fertilizer. Thus, the enrichment of dairy manure with U or AN for application on stubble-covered soil should not be recommended. Furthermore, assessments of ammonia emission from urea on crop residues should be conducted for more than seven days.

This work contributes to a better understanding of the losses of ammonia from manure and manure amended with mineral nitrogen fertilizers applied on the stubble. More studies regarding raw manure/mineral fertilizer combinations and management strategies, as acidification, that may reduce ammonia emissions from organic-mineral fertilizers are required to allow efficient use of this fertilizing strategy in no-tillage agriculture.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/IECAG2021-10017/s1>.

Author Contributions: Conceptualization, A.A.S. and D.F.; methodology, A.A.S. and D.F.; validation, D.F.; formal analysis, A.A.S. and D.F.; investigation, A.A.S. and D.F.; resources, D.F.; data curation, A.A.S.; writing—original draft preparation, A.A.S. and D.F.; writing—review and editing A.A.S. and D.F.; visualization, A.A.S. and D.F.; supervision, D.F.; project administration, D.F.; and funding acquisition, D.F. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by (1) the Project Nutri2Cycle: H2020-SFS-30-2017-“Transition towards a more carbon and nutrient efficient agriculture in Europe”, funded from the European Union, Program Horizon 2020 (Grant Agreement No 773682); (2) the Project CleanSlurry PTDC/ASP-SOL/28769/2017, “Animal slurry hygienization for use in industrial horticulture” funded by Fundação para a Ciência e Tecnologia (FCT), (3) LEAF (Linking Landscape, Environment, Agriculture and Food Research Unit), funded by FCT (UID/AGR/04129/2020). This document reflects only the authors’ view and the Union is not liable for any use that may be made of the information contained therein.

Institutional Review Board Statement: Not applicable.

Acknowledgments: The authors thank to Instituto Federal de Educação, Ciência e Tecnologia de São Paulo for supporting the PhD course of the first author.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Roy, R.N.; Finck, A.; Blair, G.J.; Tandon, H.L.S. *Plant Nutrition for Food Security—A Guide for Integrated Nutrient Management, Fertilizer and Plant Nutrition Bulletin 16*; FAO: Rome, Italy, 2006; p. 368.
2. Bouwman, L.; Goldewijk, K.K.; Van Der Hoek, K.W.; Beusen, A.H.W.; Van Vuuren, D.P.; Willems, J.; Rufino, M.C.; Stehfest, E. Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900–2050 period. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 20882–20887. [[CrossRef](#)] [[PubMed](#)]
3. Ozlu, E.; Sandhu, S.S.; Kumar, S.; Arriaga, F.J. Soil health indicators impacted by long-term cattle manure and inorganic fertilizer application in a corn-soybean rotation of South Dakota. *Sci. Rep.* **2019**, *9*, 11776. [[CrossRef](#)] [[PubMed](#)]
4. Fangueiro, D.; Alvarenga, P.; Fragoso, R. Horticulture and Orchards as New Markets for Manure Valorisation with Less Environmental Impacts. *Sustainability* **2021**, *13*, 1436. [[CrossRef](#)]
5. McConnell, D.A.; Doody, D.G.; Elliott, C.T.; Matthews, D.I.; Ferris, C.P. Impact of slurry application method on phosphorus loss in runoff from grassland soils during periods of high soil moisture content. *Ir. J. Agric. Food Res.* **2016**, *55*, 36–46. [[CrossRef](#)]
6. Iqbal, J.; Schmidt, A.M. *Tips for Winter Manure Application*; UNL: Lincoln, NE, USA, 2020; Available online: <https://cropwatch.unl.edu/2020/tips-winter-manure-application> (accessed on 5 March 2021).
7. European Commission. Collection and Analysis of Data for the Control of Emissions from the Spreading of Manure. 2014. Available online: <https://ec.europa.eu/environment/air/pdf/Final%20Report.pdf> (accessed on 4 March 2021).
8. Hassouna, M.; Eglin, T. *Measuring Emissions from Livestock Farming: Greenhouse Gases, Ammonia and Nitrogen Oxides*; INRA-ADEME Edition: Paris, France, 2016; p. 220.
9. Sommer, S.G.; Webb, J.; Hutchings, N.D. New Emission Factors for Calculation of Ammonia Volatilization from European Livestock Manure Management Systems. *Front. Sustain. Food Syst.* **2019**, *101*, 1–9. [[CrossRef](#)]
10. Fangueiro, D.; Pereira, J.L.S.; Fraga, I.; Surgy, S.; Vasconcelos, E.; Coutinho, J. Band application of acidified slurry as an alternative to slurry injection in a Mediterranean double cropping system: Agronomic effect and gaseous emissions. *Agric. Ecosyst. Environ.* **2018**, *267*, 87–99. [[CrossRef](#)]
11. Houba, V.J.G.; Temminghoff, E.J.M.; Gaikhorst, G.A.; van Vark, W. Soil analysis procedures using 0.01 M calcium chloride as extraction reagent. *Commun. Soil Sci. Plant Anal.* **2000**, *31*, 1299–1396. [[CrossRef](#)]
12. Bouwman, A.F.; Boumans, L.J.M.; Batjes, N.H. Estimation of global NH₃ volatilization loss from synthetic fertilizers and animal manure applied to arable lands and grasslands. *Glob. Biogeochem. Cycles* **2002**, *16*, 1–11. [[CrossRef](#)]

13. Feilberg, A.; Sommer, S.G. Ammonia and Malodorous Gases: Sources and Abatement Technologies. In *Animal Manure: Recycling, Treatment and Management*, 1st ed.; Sommer, S.G., Christensen, M.L., Schmidt, T., Jensen, L.S., Eds.; Wiley: West Sussex, UK, 2013; pp. 153–176.
14. Sommer, S.G.; Générumont, S.; Cellier, P.; Hutchings, N.J.; Olesen, J.E.; Morvan, T. Processes controlling ammonia emission from livestock slurry in the field. *Eur. J. Agron.* **2003**, *19*, 465–486. [[CrossRef](#)]
15. Fangueiro, D.; Hjorth, M.; Gioelli, F. Acidification of animal slurry—A review. *J. Environ. Manag.* **2015**, *149*, 46–56. [[CrossRef](#)]
16. Sigurdarson, J.J.; Svane, S.; Karring, H. The molecular processes of urea hydrolysis in relation to ammonia emissions from agriculture. *Rev. Environ. Sci. Biotechnol.* **2018**, *17*, 241–258. [[CrossRef](#)]
17. Forrestal, P.J.; Harty, M.; Carolan, R.; Lanigan, G.J.; Watson, C.J.; Laughlin, R.J.; McNeill, G.; Chambers, B.J.; Richards, K.G. Ammonia emissions from urea, stabilized urea and calcium ammonium nitrate: Insights into loss abatement in temperate grassland. *Soil Use Manag.* **2016**, *32*, 92–100. [[CrossRef](#)]
18. UNECE. *Guidance Document on Preventing and Abating Ammonia Emissions from Agricultural Source*; UN: Geneva, Switzerland, 2014; p. 100.
19. Gonzatto, R.; Carvalho Miola, E.C.; Doneda, A.; Pujol Barbosa, S.; Aita, C.; Giacomini, S.J. Volatilização de amônia e emissão de óxido nitroso após aplicação de dejetos líquidos de suínos em solo cultivado com milho. *Ciênc. Rural* **2013**, *43*, 1590–1596. [[CrossRef](#)]
20. Rochette, P.; Angers, D.A.; Chantigny, M.H.; MacDonald, J.D.; Gasser, M.O.; Bertrand, N. Reducing ammonia volatilization in a no-till soil by incorporating urea and pig slurry in shallow bands. *Nutr. Cycl. Agroecosyst.* **2009**, *84*, 71–80. [[CrossRef](#)]
21. Pujol, S.B. Emissão de Amônia e Dinâmica do Nitrogênio no Solo com Parcelamento da Dose e Adição de Inibidor de Nitrificação em Dejetos de Suínos. Doctoral Thesis, UFSM, Santa Maria, Brazil, 27 April 2012.